CONCUSSION AWARENESS

EDUCATIONAL MATERIAL ACKNOWLEDGEMENT FORM

| By my name and signature below, I acknow | wledge in accordance with Public Acts 342 and 343 of 2012 |
|---|---|
| that I have received and reviewed the Con | cussion Fact Sheet for Parents and/or the Concussion Fact |
| Sheet for Students provided by Ubly | |
| | Sponsoring Organization |
| · | |
| Participant Name Printed | Parent or Guardian Name Printed |
| | · • |
| Participant Name Signature | Parent or Guardian Name Printed |
| ¥ * | |
| Date | Date |

Return this signed form to the sponsoring organization that must keep on file for the duration of participation or age 18.

Participants and parents please review and keep the educational materials available for future reference.

Educational Material for Parents and Students (Content Meets MDCH Requirements)

Sources Michigan Department of Community Health. CDC and the National Operating Committee on Standards for Albletic Equipment (NOCSAE)

UNDERSTANDING CONCUSSION

Some Common Symptoms

Headache Pressure in the Head Nausea/Vomiting Dizziness Balance Problems
Double Vision
Blurry Vision
Sensitive to Light

Sensitive toNoise Sluggishness Haziness Fogginess Grogginess Poor Concentration Memory Problems Confusion "Feeling Down" Not "Feeling Right" Feeling Irritable Slow Reaction Time Sleep Problems

WHAT IS A CONCUSSION?

A concussion is a type of traumaticbrain injury that changes the way the brainnormally works. A concussion is caused by a fall, bump, blow or jol! to the head or bodythat causes the head and brain to movequickly back and forth. A concussion can be caused by a shaking, spiniting or a sudden stopping and starting of the head. Even a "ding," getting your bell rung," or what seems to be a mild bump or blow to the head can be serious. A concussion can happen even if you haven't been knocked out.

You can't see a concussion. Signs and symptoms of concussions can show upright after the injury or may not appear or be noticed until clays or weeks after the injury. If the student reports any symptoms of a concussion, or if yound on the symptoms yourself, seek medical attention right away. A student who may have had a concussion should not return to play on the day of the injury and until a health care professional says they are okay to return to play.

IF YOU SUSPECT A CONCUSSION:

- 1. SEEK MEDICAL ATTENTION RIGHT AWAY A health care professional will be able to decide how serious the concussionis and when it is safe for the student to return to regular activities, including sports. Don't hide it, report it. Ignoring symptoms and trying to "tough it out" often makes it worse.
- 2. KEEP YOUR STUDENT OUT OF PLAY Concussions take time to heal. Don't letthe student return to play the day of injuryand until a heath care professional says it'sokay. A student who returns to play too soon, while the brain is still healing, risks a greaterchance of having a second concussion. Young children and teens are more likely to get a concussion and take longer to recover than adults. Repeat or second concussions increase the time it takes to recover and can be very serious. They can cause permanent braindamage, aftecting the student for a lifetime. They can be fatal. It is better to miss one game than the whole season.
- 3. TELL THE SCHOOL ABOUT ANYPREVIOUS CONCUSSION-Schools should know if a student had aprevious concussion. A student's schoolmay not know about a concussion received in another sport or activity unless you notify them.

SIGNS OBSERVED BY PARENTS:

- /-ppears dażed or stunned
- Is confused about assignment or position
- Forgets an instruction
- Can't recall events prior to or after a hit
 cafell
- · Is unsure of game, score, or opponent
- · Moves clumsily

- Answers questions slowly
- Loses consciousness (even briefly)
- Shows mood, behavior, or personality changes

CONCUSSION DANGER SIGNS:

In rare cases, a dangerous blood clot may form on the brain in a person with a concussion and crowd the brain against the skull. A student should receive immediate medical attention if after a bump, blow, or jolt to the head or body s/he exhibits any of the following danger signs:

- · One pupil larger than the other
- · Is drowsy or cannot be awakened
- · A headache that gets worse
- Weakness, numbness, or decreasedcoordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Cannot recognize people/places
- Becomes increasingly confused, restlessor agitated
- Has unusual behavior
- Loses consciousness (even abriel loss of consciousness should be taken seriously.)

HOW TO RESPOND TO A REPORT OF A CONCUSSION:

If a student reports one or more symptoms of a concussion after a bump, blow, or jolt to the head or body, s/he should be kept out of athletic play the day of the injury. The student should only return to play with permission from a health care professional experienced in evaluating for concussion. During recovery, rest is key. Exercising or activities that involve a lot of concentration (such as studying, working on the computer, or playing video games) may cause concussion symptoms to reappear or get worse. Students who return to school after a concussion may need to spend fewer hours at school, take rests breaks, be given extra help and time, spend less time reading, writing or on a computer. After a concussion, returning to sports and school is a gradual process that should be monitored by a health care professional.

Remember: Concussion affects people differently. While most students with a concussion recover quickly and fully, some will have symptoms that last for days, or even weeks. A more serious concussion can last for months or longer.

To learn more, go to www.cdc.gov/concussion.

Parents and Students Must Sign and Return the Educational Material Acknowledgement Form